



Key impacts of climate engineering on biodiversity and ecosystems, with priorities for future research

Caitlin G. McCormack, Wanda Born, Peter J. Irvine, Eric P. Achterberg, Tatsuya Amano, Jeff Ardron, Pru N. Foster, Jean-Pierre Gattuso, Stephen J. Hawkins, Erica Hendy, W. Daniel Kissling, Salvador E. Lluch-Cota, Eugene J. Murphy, Nick Ostle, Nicholas J.P. Owens, R. Ian Perry, Hans O. Pörtner, Robert J. Scholes, Frank M. Schurr, Oliver Schweiger, Josef Settele, Rebecca K. Smith, Sarah Smith, Jill Thompson, Derek P. Tittensor, Mark van Kleunen, Chris Vivian, Katrin Vohland, Rachel Warren, Andrew R. Watkinson, Steve Widdicombe, Phillip Williamson, Emma Woods, Jason J. Blackstock & William J. Sutherland

To cite this article: Caitlin G. McCormack, Wanda Born, Peter J. Irvine, Eric P. Achterberg, Tatsuya Amano, Jeff Ardron, Pru N. Foster, Jean-Pierre Gattuso, Stephen J. Hawkins, Erica Hendy, W. Daniel Kissling, Salvador E. Lluch-Cota, Eugene J. Murphy, Nick Ostle, Nicholas J.P. Owens, R. Ian Perry, Hans O. Pörtner, Robert J. Scholes, Frank M. Schurr, Oliver Schweiger, Josef Settele, Rebecca K. Smith, Sarah Smith, Jill Thompson, Derek P. Tittensor, Mark van Kleunen, Chris Vivian, Katrin Vohland, Rachel Warren, Andrew R. Watkinson, Steve Widdicombe, Phillip Williamson, Emma Woods, Jason J. Blackstock & William J. Sutherland (2016) Key impacts of climate engineering on biodiversity and ecosystems, with priorities for future research, Journal of Integrative Environmental Sciences, 13:2-4, 103-128, DOI: [10.1080/1943815X.2016.1159578](https://doi.org/10.1080/1943815X.2016.1159578)

To link to this article: <https://doi.org/10.1080/1943815X.2016.1159578>



View supplementary material [↗](#)



Published online: 23 Mar 2016.



Submit your article to this journal [↗](#)



Article views: 1333



View related articles [↗](#)



View Crossmark data [↗](#)

Key impacts of climate engineering on biodiversity and ecosystems, with priorities for future research

Caitlin G. McCormack^a, Wanda Born^b, Peter J. Irvine^b, Eric P. Achterberg^{c,d}, Tatsuya Amano^a, Jeff Ardrone^e, Pru N. Foster^f, Jean-Pierre Gattuso^{g,h}, Stephen J. Hawkins^c, Erica Hendy^{f,i}, W. Daniel Kissling^j, Salvador E. Lluch-Cota^k, Eugene J. Murphy^l, Nick Ostle^m, Nicholas J.P. Owensⁿ, R. Ian Perry^o, Hans O. Pörtner^p, Robert J. Scholes^q, Frank M. Schurr^r, Oliver Schweiger^s, Josef Settele^{s,t}, Rebecca K. Smith^a, Sarah Smith^u, Jill Thompson^v, Derek P. Tittensor^{u,w}, Mark van Kleunen^x, Chris Vivian^y, Katrin Vohland^z, Rachel Warren^{aa}, Andrew R. Watkinson^{aa}, Steve Widdicombe^{ab}, Phillip Williamson^{ac}, Emma Woods^{ad}, Jason J. Blackstock^{ae} and William J. Sutherland^a

^aConservation Science Group, Department of Zoology, University of Cambridge, Cambridge, UK; ^bSustainable Interactions with the Atmosphere, Institute for Advanced Sustainability Studies e.V., Potsdam, Germany; ^cOcean and Earth Science, National Oceanography Centre Southampton, University of Southampton Waterfront Campus, Southampton, UK; ^dGEOMAR, Helmholtz Centre for Ocean Research, Kiel, Germany; ^eGlobal Contract for Sustainability, Institute for Advanced Sustainability Studies e.V., Potsdam, Germany; ^fSchool of Earth Sciences, University of Bristol, Bristol, UK; ^gSorbonne Universités, UPMC, Univ Paris 06, CNRS-INSU, Laboratoire d'Océanographie de Villefranche, Villefranche-sur-Mer, France; ^hInstitute for Sustainable Development and International Relations, Sciences Po, Paris, France; ⁱSchool of Biological Sciences, University of Bristol, Bristol, UK; ^jInstitute for Biodiversity and Ecosystem Dynamics (IBED), University of Amsterdam, Amsterdam, The Netherlands; ^kPrograma de Ecología Pesquera, Centro de Investigaciones Biológicas del Noroeste (CIBNOR), La Paz, Mexico; ^lBritish Antarctic Survey, Cambridge, UK; ^mLancaster Environment Centre, Lancaster University, Lancaster, UK; ⁿScottish Association for Marine Science, Scottish Marine Institute, Oban, UK; ^oDepartment of Fisheries and Oceans, Pacific Biological Station, Nanaimo, Canada; ^pAlfred-Wegener-Institut für Polar- und Meeresforschung, Ökophysiologie, Germany; ^qCouncil for Scientific and Industrial Research, Pretoria, South Africa; ^rInstitut des Sciences de l'Évolution de Montpellier, UMR-CNRS 5554, Université Montpellier II, Montpellier, France; ^sDepartment of Community Ecology, UFZ Centre for Environmental Research, Halle, Germany; ^tiDiv, German Centre for Integrative Biodiversity Research (iDiv) Halle-Jena-Leipzig, Leipzig, Germany; ^uUNEP World Conservation Monitoring Centre, Cambridge, UK; ^vCentre for Ecology and Hydrology, Midlothian, UK; ^wDepartment of Biology, Dalhousie University, Halifax, Canada; ^xEcology, Department of Biology, University of Konstanz, Konstanz, Germany; ^yCefas, Lowestoft Laboratory, Lowestoft, UK; ^zMuseum für Naturkunde, Leibniz-Institut für Evolutions- und Biodiversitätsforschung, Berlin, Germany; ^{aa}School of Environmental Sciences, University of East Anglia, Norwich, UK; ^{ab}Plymouth Marine Laboratory, Plymouth, UK; ^{ac}Natural Environment Research Council and School of Environmental Sciences, University of East Anglia, Norwich, UK; ^{ad}The Royal Society, London, UK; ^{ae}Science, Technology, Engineering and Public Policy, University College London, London, UK

ABSTRACT

Climate change has significant implications for biodiversity and ecosystems. With slow progress towards reducing greenhouse gas emissions, climate engineering (or 'geoengineering') is receiving increasing attention for its potential to limit anthropogenic climate change and its damaging effects. Proposed techniques, such as ocean fertilization for carbon dioxide removal or stratospheric sulfate

ARTICLE HISTORY


Received 12 July 2015

Accepted 25 February 2016

KEYWORDS

Biodiversity; carbon dioxide removal; climate engineering; ecosystems; geoengineering; solar radiation management

CONTACT William J. Sutherland  w.sutherland@zoo.cam.ac.uk

 Supplemental data for this article can be accessed here. [<http://dx.doi.org/10.1080/1943815X.2016.1159578>]

injections to reduce incoming solar radiation, would significantly alter atmospheric, terrestrial and marine environments, yet potential side-effects of their implementation for ecosystems and biodiversity have received little attention. A literature review was carried out to identify details of the potential ecological effects of climate engineering techniques. A group of biodiversity and environmental change researchers then employed a modified Delphi expert consultation technique to evaluate this evidence and prioritize the effects based on the relative importance of, and scientific understanding about, their biodiversity and ecosystem consequences. The key issues and knowledge gaps are used to shape a discussion of the biodiversity and ecosystem implications of climate engineering, including novel climatic conditions, alterations to marine systems and substantial terrestrial habitat change. This review highlights several current research priorities in which the climate engineering context is crucial to consider, as well as identifying some novel topics for ecological investigation.

1. Introduction

Anthropogenic emissions of greenhouse gases including carbon dioxide are considered the main cause of an observed 0.8 °C increase in average global surface temperature since pre-industrial times (IPCC 2013). These changes in greenhouse gas concentrations have implications not only for temperature, but also for precipitation, ice-sheet dynamics, sea levels, ocean acidification and extreme weather events (IPCC 2013). Such changes are already starting to have substantive effects on biodiversity and ecosystems, including altered species' distributions, interspecific relationships and life history events, and are predicted to intensify into the future (Chen et al. 2011; Bellard et al. 2012; Warren et al. 2013). With continued high greenhouse gas emissions (Jackson et al. 2016; International Energy Agency 2015), climate engineering ('geoengineering') has been receiving increasing attention for its potential to be used to counteract climate change and reduce its damaging effects (IPCC 2013).

Climate engineering refers to large-scale interventions in the Earth system intended to counteract climate change. There are two main types (see Figure 1, Table 1 and Supporting Information1 in Supporting Information): (a) carbon dioxide removal (CDR) techniques, designed to reduce atmospheric carbon dioxide concentrations, and (b) solar radiation management (SRM), designed to reflect solar radiation away from Earth (The Royal Society 2009; Secretariat of the Convention on Biological Diversity 2012; Caldeira et al. 2013). There are a range of other terms for these processes. If effective the primary impact of climate engineering would be to reduce the damaging effects of climate change; CDR by reducing CO₂ concentrations to abate the process of climate change itself and SRM by direct lowering of global temperatures. All techniques will also have secondary impacts associated with their implementation, ranging from local land-use changes to globally reduced stratospheric ozone levels, for example (Ricke et al. 2010; Secretariat of the Convention on Biological Diversity 2012; Tilmes et al. 2013). These secondary impacts have wide-reaching and potentially complex biodiversity implications (Winder 2004). However, the possible consequences and the research needed to determine them, have received little attention from the ecological research community and are largely absent from climate engineering discussions (Russell et al. 2012).




Figure 1. Schematic of climate engineering techniques considered in this review, covering CDR techniques and SRM techniques.

The current lack of consideration of climate engineering impacts on biodiversity and ecosystems is due in part to the number, complexity, novelty, and large spatial and temporal scale of the potential effects. It is difficult or impossible to empirically test the effects of most of the techniques (Keith 2000; MacMynowski et al. 2011; Keller et al. 2014) and deciding on the most pressing research topic can be challenging. The issue can seem an overwhelming challenge for ecological science, causing research to respond slowly, and to follow rather than inform policy decisions (Sutherland & Woodroof 2009). Climate engineering has already entered policy discussions (Secretariat of the Convention on Biological Diversity 2012; International Maritime Organization 2013; IPCC 2013) and, to date, although implementation is regulated, there is no comprehensive international agreement covering all climate engineering techniques (Rickels et al. 2011). It is therefore critical that research to understand potential ecological effects of climate engineering begins as soon as possible so that it can inform the development of ecologically-sensitive techniques and evidence-based policy decisions.

For this study, a process of literature review and expert consultation was used to review the potential biodiversity and ecosystem effects of climate engineering. We focus on the potential side-effects of implementing the techniques rather than the anticipated climate change amelioration effect as the former have received relatively little attention and the latter is a large and complex body of ongoing research beyond the scope of the current project. We identify key areas where climate engineering presents important questions that should be considered within existing priority ecological research efforts, as well as identifying a number of novel knowledge gaps. We suggest a list of research questions which we hope will encourage timely investigation of the potential ecological effects of climate engineering.

2. Materials and methods

'Horizon-scanning' involves the systematic assessment of emerging threats and opportunities, in order to identify key upcoming issues (Sutherland 2006; Sutherland & Woodroof 2009; Martin et al. 2012; Sutherland et al. 2012). In the current study, an adapted process called 'impact scanning' was used; impacts of climate engineering were identified from the literature and reviewed to prioritize those which are likely to have the greatest effects on

biodiversity and ecosystems. The degree of scientific understanding about the effects was also evaluated, to identify critical knowledge gaps. An expert consultation process combining elements of the Nominal Group and Delphi techniques (Hutchings & Raine 2006) was used (Figure 2 gives a summary). Participants gave verbal consent to take part in this exercise. We did not obtain formal written consent as all data and comments are kept anonymous and it was agreed from the outset that participants were to be authors of the resulting paper and approve its contents prior to publication.

2.1. Literature reviews

A literature review was conducted to identify the potential biodiversity and ecosystem effects of climate engineering techniques. As the scope of the existing literature was uncertain, the recent reports of the Royal Society (2009) and the Secretariat of the Convention on Biological Diversity (2012) were used as a starting point. An approach based on snowball sampling (Biernacki & Waldorf 1981) was used to identify further relevant literature from




Figure 2. Flow diagram of study methodology.

their citations, and then from the citations of these citations, and so on. Seventeen geoengineering techniques were included in the review (Figure 1) based on those discussed in prominent literature at the time (The Royal Society 2009; Rickels et al. 2011). Overall, the review found 154 environmental changes predicted to result from the techniques, each with a range of associated potential biodiversity and ecosystem effects (Supporting Information S1). Additional environmental changes were added by the participating group of researchers so that a total of 192 changes and their associated effects were assessed in total. The focus was on the side-effects of the implementation of the techniques, rather than the effects they would cause by counteracting climate change, which is beyond the scope of the current study. In a separate literature review, assessments of the technical feasibility and anticipated effectiveness of the techniques were identified using the same literature sampling technique as above, and used to shortlist five techniques about which research questions were formulated.

2.2. Scoring round 1: survey

The assessment was conducted by a working group of 34 senior academic scientists with expertise in biodiversity, ecosystems and environmental and climatic change. Participants were identified through internet searches and selected to ensure an even split between terrestrial and marine expertise, and a global scope; the majority of experts were based at European institutions but there were also representatives from Canada, North America, Mexico and South Africa, and all had extensive knowledge of ecosystems beyond their institution's country.

Each participant first completed an Excel-based survey exercise. They read the report of the literature review of biodiversity and ecosystem effects of climate engineering (Supporting Information S1), and used the information to score a list of environmental changes for each of the techniques between 0 and 100, to reflect the relative importance of their potential effects on biodiversity and ecosystems. They added comments to explain their scores. Each climate engineering technique was considered separately. At the end of the survey, the participants compared their top prioritised environmental changes from each technique and scored them between 0 and 100. These values were used as 'swing weights' to calibrate the earlier scores, making them comparable across the techniques (Holt 1996). In a second Excel-based survey, participants used the literature review report in combination with their own experience and expertise to score the environmental changes between 0 and 100 to reflect the extent of scientific knowledge about their biodiversity and ecosystem effects. They also suggested priority research questions. Detailed guidelines and definitions were provided for both survey exercises to ensure that scores were comparable amongst participants. They were asked to assume deployment of the technique at a 'climatically-significant scale' (Lenton & Vaughan 2009; Williamson et al. 2012) and against a background of climate change causing a warming world with an acidifying ocean. SRM-induced climate changes were considered independently of the concurrent greenhouse gas-induced climate changes. Nevertheless, the biodiversity and ecosystem consequences identified are equally applicable when the two drivers are considered together.

2.3. Re-scoring

A summary of the survey responses was sent to each expert for them to review ahead of a two day workshop in May 2013. At the workshop, participants shared reasons for their scores, and heard perspectives from others in the group. Parallel groups discussed a subset of the climate engineering techniques and their associated environmental changes and biodiversity and ecosystem effects. Following discussion, the experts then individually re-scored using the same 0–100 scale or kept their original score based on the discussion.

In a final session, the research questions suggested during the second pre-workshop survey were reviewed and refined.

2.4. Calculating an ‘index of priority’

A median was calculated from the group’s final importance and scientific understanding scores (both using range of 0–100). This was used to calculate an ‘index of priority’ for each of the environmental changes across all of the climate engineering techniques, using the equation: $(\text{Importance score} + (100 - \text{Understanding score})) \times 0.5$.

The index of priority was used to rank the environmental changes; a change is of greater priority if it has more important potential effects on biodiversity and ecosystems and/or there is less understanding about its effects. A list of the top 20 changes across all of the techniques was identified from the results of this scoring.

2.5. Shortlisted techniques and research questions

As well as assessing the effects across all 17 climate engineering techniques, we specifically assessed effects associated with techniques that we concluded were more plausible for implementation than others; five of the 17 climate engineering techniques were identified from a review of existing assessments as having relatively higher anticipated efficacy (potential climate change forcing when deployed at maximum scale) and technical feasibility (availability of materials, technology and knowledge to implement) than the other techniques (Table 1) (e.g. (Lenton & Vaughan 2009; The Royal Society 2009; Caldeira et al. 2013). This was taken to indicate that they are more plausible options for implementation, meaning that potential effects associated with them are the most pertinent to consider.

The index of priority was used to identify two or three highest priority environmental changes associated with each of these five techniques. The expert group identified key knowledge gaps and research questions about the potential biodiversity and ecosystem effects, using the questions suggested during the survey as a starting point.

3. Results and discussion

3.1. Key themes for research – across all techniques

The ‘index of priority’ was used to first rank all of the environmental changes across all of the 17 climate engineering techniques, assuming equal likelihood of implementation. A full list of the median scores and index of priority values is given in Supporting Information S4. The top 20 of these environmental changes (Table 2), and patterns within the rest of the ranked list, reveals interesting themes in the types of changes that were judged by

Table 1. Description of climate engineering techniques and shortlisting on the basis of technical feasibility, affordability and/or anticipated effectiveness.

Climate engineering technique	SRM or CDR	Description	Prioritization	Reasons for prioritization
<i>High priority techniques</i>				
Ocean fertilization – iron	CDR	Soluble iron minerals added to regions of the ocean where availability limits productivity. Cover c. 30% of the ocean surface, including the Southern Ocean, and the equatorial and northern Pacific ^a	High	Field experimentation ^b shows enhanced CO ₂ uptake can be achieved. Iron has greater potential CO ₂ sequestration per amount of nutrient added compared to macronutrient fertilization ^b so is prioritized over nitrogen/phosphorus (below)
Bio-energy with carbon capture and storage (BECCS)	CDR	Biomass burned for fuel and CO ₂ emissions produced during processing and combustion captured and transferred to long-term geological or ocean storage ^{a,c}	High	Techniques for bioenergy production, processing, combustion, and capture and storage of CO ₂ already developed ^{a,c} . Relatively high anticipated CO ₂ sequestration potential ^{b,d,e}
Marine cloud albedo	SRM	Reflectivity of clouds over the ocean is enhanced by increasing the number of particles which act as cloud condensation nuclei, by spraying seawater into clouds ^{a,e}	High	Potential for large radiative forcing effect. ^{e,f} Potentially technically feasible and relatively affordable technology ^{a,g,h}
Stratospheric sulfate aerosols	SRM	Sulfur dioxide or hydrogen sulfide injected into the lower stratosphere to form sulfate aerosol particles which scatter incoming shortwave radiation ^d	High	Potential for large radiative forcing effect. ^{e,f} Potentially technically feasible and relatively affordable technology ^d
Direct air capture (DAC)	CDR	Free-standing structures constructed in areas with good airflow. Sorbent materials on surfaces selectively trap CO ₂ from ambient air. Isolated CO ₂ transferred to a long-term geological or ocean store ^a	High	High anticipated CO ₂ sequestration potential. ^{e,f} Relatively achievable technological requirements ^a
<i>Lower priority techniques</i>				
Ocean fertilization – nitrogen/phosphorus	CDR	Soluble phosphorus or nitrogen minerals added to regions of the ocean where availability limits productivity. These regions cover 40% of the ocean surface including tropical and subtropical gyres ^{a,b}	Low	Limited carbon sequestration potential ^{b,f} . Significant volumes of mined minerals required ^a
Biomass – storage in the ocean	CDR	Terrestrial biomass harvested, baled and deposited onto the sea floor below 1000–1500 m where conditions limit decomposition ^{a,i}	Low	Unlikely to be viable at a scale to appreciably offset global CO ₂ emissions. ^a Requires novel techniques and equipment
Biochar	CDR	Biomass burned in low oxygen ('pyrolysis') to form solid product similar to charcoal. This is dug into soils where it acts as a carbon reservoir ^{a,i}	Low	Feasibility and anticipated effectiveness in achieving net CO ₂ reduction limited by significant land use requirements ^{a,i}
Enhanced weathering <i>in situ</i>	CDR	CO ₂ dissolved in solution and injected into basic rocks in the Earth's crust to react with basic minerals such as olivine to form mineral compounds ^a	Low	Significant logistical challenges and uncertainty over chemical feasibility and energy requirements ^a
Afforestation or reforestation	CDR	Forest established on currently non-forested land to increase CO ₂ uptake and storage through photosynthesis ^{a,i}	Low	Biodiversity and ecosystem effects of afforestation and reforestation have previously been subject to detailed reviews so are not considered here
Enhanced weathering: to land	CDR	Basic rock minerals – such as olivine – are quarried, ground into fine particles and spread on soils to undergo accelerated weathering, reacting with atmospheric CO ₂ and converting it to mineral compounds ^{i,k}	Low	Relatively good technical feasibility but high energy requirements and CO ₂ emissions associated with quarrying, processing and spreading materials ^{a,i,k}

(Continued)

Table 1. (Continued)

Climate engineering technique	SRM or CDR	Description	Prioritization	Reasons for prioritization
Enhanced weathering: to ocean	CDR	Quarried and processed carbonate or silicate materials are added to the surface ocean. The basic/alkaline materials react with CO ₂ in the water, converting it to bicarbonate ions. CO ₂ content of the ocean is reduced allowing more to be absorbed from the atmosphere ⁱ	Low	[See. Enhanced weathering: to land]
Enhanced upwelling/downwelling	CDR	The natural process of upwelling – deep-ocean waters brought to the surface by ocean circulation – is enhanced using man-made pipes and pumps. Water brought to the surface is rich in nutrients and cooler than existing surface waters, leading to increased uptake of atmospheric CO ₂ . Alternatively, natural downwelling would be enhanced by cooling CO ₂ -rich ocean surface waters, causing them to sink to the deep ocean ^{aj}	Low	Very limited potential to achieve net drawdown of CO ₂ due to high CO ₂ content of waters brought to surface by both techniques. ^b Significant logistical and engineering challenges ⁱ
Surface albedo – urban	SRM	Albedo of urban structures increased using bright paint or materials ^{sa,m}	Low	Very low anticipated radiative forcing potential and therefore low cost-effectiveness ^{a,e,f}
Surface albedo – desert	SRM	Albedo of desert regions – which receive a high proportion of incoming solar radiation – increased by covering areas in man-made reflective materials ^{sa,f}	Low	Very low anticipated affordability and very large land requirements ^a
Surface albedo – crop	SRM	Plants selected for high surface albedo are established over large areas of cropland or grassland/shrubland ^{a,m,n}	Low	Low anticipated radiative forcing potential, ^{d,e,f} scale of implementation required for measurable effect prohibitively large ^{e,f}
Sunshades	SRM	Sun shields or deflectors are installed in space to reflect a proportion of sunlight away from the Earth ^{ad}	Low	Very low timeliness and affordability ^{a,d}

^aThe Royal Society 2009.
^bWilliamson et al. 2012.
^cMetz et al. 2005.
^dCaldeira et al. 2013.
^eLenton & Vaughan 2009.
^fVaughan & Lenton 2011.
^gFoster et al. 2013.
^hLatham et al. 2012.
ⁱSecretariat of the Convention on Biological Diversity 2012.
^jMatthews et al. 2002.
^kHartmann et al. 2013.
^lZhou & Flynn 2005.
^mIrvine et al. 2011.
ⁿSingarayer et al. 2009.

Table 2. Top environmental changes across all techniques presented in rank order according to an 'index of priority'. A higher value indicates a greater priority for research due to higher judged importance and/or lower scientific understanding of potential biodiversity and ecosystem effects. See Supporting Information S4 for a full list of environmental changes and scores.

Rank	Technique	SRM or CDR	Environmental change	Median importance score (interquartile range) 100 = highest importance	Median scientific understanding score (interquartile range) 0 = no scientific understanding; 100 = complete scientific understanding	Index of priority* (100 = highest priority)
1	Solar radiation management 'dimming' techniques ^a	SRM	The 'termination effect' ^b : Rapid increase of global temperatures if solar radiation management failed or was terminated	99.9 (6)	20 (5)	90
2	Solar radiation management 'dimming' techniques ^a	SRM	Regionally-variable changes in precipitation due to altered atmospheric circulation. Increase in some areas, decrease in others	80 (18)	30 (10)	75
3	Solar radiation management 'dimming' techniques ^a	SRM	Creation of high CO ₂ /low temperature climate (unlike either the current low CO ₂ /low temperature conditions or high CO ₂ /high temperature conditions of projected climate change)	70 (27)	20 (8)	75
4	Solar radiation management 'dimming' techniques ^a	SRM	Reduced amplitude of seasonal temperature range with warmer winters and cooler summers	75 (20)	30 (10)	73
5	Solar radiation management 'dimming' techniques ^a	SRM	Small but detectable global cooling within ~5 years of solar radiation management deployment (relative to elevated temperatures caused by global warming effect)	74 (11)	30 (5)	72
6	Solar radiation management 'dimming' techniques ^a	SRM	Reduced equator-to-pole temperature gradient due to greater reduction in incoming solar radiation at the tropics than at higher latitudes	70 (19)	30 (6)	70
7	Solar radiation management 'dimming' techniques ^a	SRM	Slowing of the global hydrological cycle (reduced evaporation and precipitation)	70 (15)	30 (10)	70
8	Enhanced desert albedo	SRM	Potentially strong reduction in continental rainfall, particularly in monsoon regions	64 (15)	30 (8)	68
9	Enhanced upwelling/downwelling	CDR	Increased primary productivity in surface ocean as a result of artificially enhanced upwelling of nutrient-rich deep waters (in mid-ocean locations)	63 (25)	30 (23)	67
10	Solar radiation management 'dimming' techniques ^a	SRM	Changes in ocean circulation patterns due to changes in energy into and out of the ocean due to reduced atmospheric temperature	63 (17)	30 (10)	67
11	Ocean fertilization with iron	CDR	Increased primary productivity in high nutrient low chlorophyll regions of the ocean due to iron fertilization	70 (30)	40 (15)	66
12	Enhanced upwelling/downwelling	CDR	Increased area of man-made structures in the ocean for artificial enhancement of upwelling or downwelling	55 (20)	25 (16)	65
13	Biomass: storage in the ocean	CDR	Increased nutrient availability in deep ocean and on sea floor due to deposition of harvested terrestrial biomass	50 (23)	15 (18)	65

(Continued)

Table 2. (Continued)

Rank	Technique	SRM or CDR	Environmental change	Median importance score (interquartile range) 100 = highest importance	Median scientific understanding score (interquartile range) 0 = no scientific understanding; 100 = complete scientific understanding	Index of priority* 100 = highest priority)
14	Enhanced cropland or grassland albedo	SRM	Establishment of monocultures of high-reflectivity vegetation over several million km ² to replace natural and semi-natural grassland and shrubland habitats	80 (17)	50 (28)	65
15	Biomass: storage in the ocean	CDR	Reduced oxygen in deep ocean due to decomposition of introduced organic matter (harvested terrestrial biomass)	55 (33)	30 (28)	65
16	Enhanced cropland or grassland albedo	SRM	Conversion of (dark) forest habitats to establish (lighter) grassland or cropland	79 (25)	50 (30)	63
17	Biomass: storage in the ocean	CDR	Large-scale coverage (smothering) of deep-ocean seabed with harvested terrestrial biomass	52 (47)	25 (15)	63
18	Enhanced weathering: base materials to land	CDR	Change in soil properties with addition of powdered basic rock (soil structure, density, aggregation and water retention)	9 (9)	30 (10)	63
19	Enhanced desert albedo	SRM	Large-scale covering of desert surface with man-made materials	50 (13)	25 (23)	61
20	Ocean fertilization: nitrogen or phosphorus	CDR	Increased primary productivity in low nutrient low chlorophyll regions of the ocean due to nitrate or phosphate fertilization	60 (20)	40 (13)	60

*The 'Index of priority' is calculated by: (Importance score + (100 – Understanding score)) × 0.5.

^aSRM 'dimming' techniques refers to sunshades, stratospheric sulfate aerosols and enhanced marine cloud albedo, which reflect a proportion of incoming solar radiation back into space. Environmental changes under this heading are taken to be common to these three techniques.

^bThe termination effect is associated with the possible failure or termination of SRM 'dimming' techniques, rather than their implementation or functioning.

the expert group to have important biodiversity and ecosystem consequences but limited scientific understanding.

3.1.1. Climatic changes

The top seven of the 20 prioritized environmental changes (Table 2) recognize the potentially substantial and complex biodiversity and ecosystem implications of global-scale alterations to climatic processes associated with SRM ‘dimming’ techniques – sunshades, sulfate aerosols and enhanced marine cloud albedo. These techniques reduce incoming shortwave radiation to the earth, reducing global mean surface temperature, but causing regionally variable changes in climatic conditions (Caldeira et al. 2013), such as potential enhancement of increases or decreases in precipitation caused by climate change (Irvine et al. 2010; Ricke et al. 2010; Kravitz, Robock, et al. 2013). ‘Novel’ regional climatic states could occur (Irvine et al. 2010). The ecological effects of these are challenging to predict (Williams et al. 2007).

Changes to temperature and precipitation patterns were considered by the group to be highly important for biodiversity and ecosystems as they are strong determinants of species’ life history, phenology, physiological performance, distribution and interactions (Pörtner & Farrell 2008; Cahill et al. 2013). A reduction in the equator-to-pole temperature gradient, for example, would shift species’ climatic ranges (Couce et al. 2013), which would lead to altered ecological community assemblages and a change in the distribution of biomes (Walther et al. 2002; Burrows et al. 2011). Changes in the amplitude of seasonal temperature variation could strongly influence the timing of ecological processes such as migration, breeding, flowering and phytoplankton blooms (Sims et al. 2001; Edwards & Richardson 2004; Menzel et al. 2006). Both the climatic effects and the biodiversity impacts they cause are likely to be highly regionally variable, due to factors such as local microclimatic conditions (De Frenne et al. 2013), or circulation patterns in the marine environment, meaning there are large gaps in knowledge and understanding of the effects and a need for research.

Changes affecting precipitation and surface water availability were also prioritized; regionally variable changes to precipitation patterns, the slowing of the global hydrological cycle (Tilmes et al. 2013), and a potential reduction in continental rainfall associated with enhanced desert albedo (Irvine et al. 2011), were all included in the top 20 (Table 2). Water availability influences rates of primary productivity and the composition of plant communities that underpin terrestrial habitats (Cleland et al. 2013). Determining the trajectory of the ecological effects of changing precipitation patterns is subject to uncertainty due to differences in individual and species responses, which compound uncertainties over the likely direction and magnitude of the precipitation change (Mustin et al. 2007; Hoffmann & Sgro 2011). Paleocological records of responses to past precipitation changes – for example, the ‘greening’ of the Sahara – can offer some indication of potential effects (e.g. Willis et al. 2013), as can ongoing research on effects of precipitation changes associated with climate change, but specific research needs to be conducted in the context of climate engineering scenarios.

3.1.2. Changes affecting marine ecosystems

Many of the prioritized environmental changes are associated with ocean systems (Table 2). Already, anthropogenic emissions of CO₂ are causing ocean acidification due to increased dissolved inorganic carbon in ocean waters. Such chemical changes have potential impacts on the acid-base balance, metabolic energy allocation and calcification of marine organisms (Bopp et al. 2013; Kroeker et al. 2013). SRM techniques would not address atmospheric

CO₂, so in the absence of additional actions to reduce greenhouse gas levels, concentrations will almost certainly increase relative to present day, which could lead to worsening acidification (Keller et al. 2014). However, there is uncertainty about the net effect; for the same emission rates, SRM could lessen CO₂ rise in the atmosphere by causing enhanced terrestrial CO₂ uptake and by avoiding positive feedbacks (e.g. carbon release from thawing tundra, fire etc.; see Matthews et al. 2009). The net effect of SRM on ocean acidification could therefore be slightly beneficial compared to a non-SRM scenario. However, SRM will also reduce sea-surface temperatures, which affect CO₂ dissolution rates, ocean circulation and other poorly-understood feedback processes, so the overall effect is uncertain (Williamson & Turley 2012). The relationship between temperature and ocean acidification impacts on marine calcifiers, and ecosystems dependent on carbonate structures (e.g. coral reefs), is an area of active research (e.g. Anthony et al. 2011) but has so far received little attention in the climate engineering context. To date, only one study (Couce et al. 2013) has investigated these potential implications of SRM, and finds that moderate deployment could reduce degradation of global coral reef habitat compared to no SRM, according to model simulations.

SRM 'dimming' techniques will affect global ocean circulation through changes to the energy exchanges between the ocean and the atmosphere (McCusker et al. 2012). Light availability (partially determined by incoming solar irradiance), temperature, and nutrient patterns fundamentally determine marine ecological communities, and are responsible for diversity both between ocean strata and across latitudes. Changes to circulation will alter these factors, with the potential for biodiversity consequences throughout the entire marine system (Drinkwater et al. 2010; Hardman-Mountford et al. 2013). The group's scores indicate there is limited scientific understanding of the likely biodiversity and ecosystem effects, particularly as they will vary regionally (Secretariat of the Convention on Biological Diversity 2012). The group acknowledged that oceanic islands would be highly vulnerable to changes in ocean-atmosphere dynamics (e.g. Loope & Giambelluca 1998). These habitats often support a high concentration of endemic species and their populations are generally small and geographically isolated, restricting their ability to adapt. Novel impacts of climate engineering could also affect them, such as possible deposition of sea water used for enhanced cloud albedo; this could further reduce freshwater availability, which is often limited on islands (Meehl 1996).

Increased primary productivity in the surface ocean due to artificially enhanced fertilization is judged to be a highly important change across the various CDR fertilization methods (Table 2). The phytoplankton communities that would be directly impacted underpin a significant proportion of ocean ecological communities and determine parameters such as light penetration, nutrient cycling, and the supply of organic material to benthic systems (Falkowski et al. 1998; Kirk 2011). Ocean fertilization could therefore have profound effects throughout marine ecosystems, particularly in currently low-productivity areas (Falkowski et al. 1998). 'Knock-on' trophic effects observed in open-ocean fisheries, whereby changes in one group of species has broad effects throughout the ecosystem (e.g. Bailey et al. 2009), would very likely occur. Effects are likely to be widely spread by global ocean circulation (Williamson et al. 2012). Although their effects are sometimes conflated in the climate engineering literature, we suggest that it is critical to distinguish iron fertilization in high nutrient low chlorophyll ocean regions from nitrogen or phosphorous fertilization in low nutrient low chlorophyll regions. Field trials of iron fertilization have shown varying impacts on phytoplankton communities and the marine ecosystem (Williamson et al. 2012) and a diversity of

effects can also be anticipated to result from nitrogen or phosphorus fertilization (Lampitt et al. 2008). Increased productivity caused by enhanced upwelling/downwelling was judged to be less well understood and so was the highest prioritized; modeling suggests that intended effects of enhanced vertical mixing may be less strong than anticipated, will vary greatly from place to place, and may even be opposite from that desired (Dutreuil et al. 2009). The engineered structures required for enhanced upwelling were also judged to have important biodiversity and ecosystem implications, creating artificial reefs or acting as 'stepping stones' for species migration, distribution, and aggregation (Mineur et al. 2012).

3.1.3. Changes affecting the deep ocean

Environmental changes with effects in the deep ocean were repeatedly identified as priorities for further research by the group (Table 2). There is a general lack of knowledge about these environments (Costello et al. 2010) but fisheries research indicates that deep sea species are sensitive to disturbance and slow to recover (e.g. Devine et al. 2006). It is therefore likely that effects of climate engineering techniques on the deep sea would be long-lasting. Large-scale coverage of the deep-ocean seabed, associated with the technique biomass storage in the ocean (Table 1), would be a significant alteration of relatively undisturbed habitats. Reduced oxygen and enhanced nutrient levels due to decaying organic matter could impact species richness, physiological processes and community composition (Levin et al. 2001; Lampitt et al. 2008). There is a need to increase fundamental understanding of these environments before deployment of any climate engineering technique that might impact them.

3.1.4. Large-scale terrestrial habitat disturbance or destruction

Large-scale disturbance of terrestrial habitats was a topic prioritized by the group, and could result from a number of climate engineering techniques (Supporting Information S1). Although the effects of such habitat change are considered to be relatively well understood (Table 2), the anticipated scale associated with climate engineering on a 'climatically significant' scale is considerable and would be additional to current processes. Specifically, the replacement of (semi-)natural grassland and shrubland, or forest habitats, with reflective plants to increase surface albedo for SRM was included in the 20 priority changes (Table 2). This conversion of existing habitat constitutes complete habitat loss for inhabitant species (Secretariat of the Convention on Biological Diversity 2012). Detrimental effects could be reduced by limiting planting to degraded land (e.g. Tilman et al. 2009). However, the area required in order for the technique to impact the global climate would inevitably exceed this resulting in conversion of natural or semi-natural habitats (see Lenton & Vaughan 2009; Tilman et al. 2009).

Alteration or loss of desert habitats through coverage with manmade reflective materials (an SRM technique) is also included within the 20 prioritized changes (Table 2). It is estimated that to offset the warming from a doubling of atmospheric CO₂ concentrations, an area of approximately 12 million square kilometers – roughly 1.2 times the area of the Saharan desert – would need to be covered (Lenton & Vaughan 2009; Vaughan & Lenton 2011). Although considered to have low biodiversity, desert regions contain many endemic species that are highly adapted to the local conditions. They are likely to be significantly affected by a long-term increase in shading and change in regional temperatures caused by man-made structures (Stahlschmidt et al. 2011). Alteration of the habitats may allow other

species to become established in desert regions, leading to changes in the unique ecological community composition (Steidl et al. 2013).

3.1.5. Alteration of soil properties

Another essential area for research was the impact of climate engineering on soils. Specifically, changes in soil properties due to the addition of powdered alkali rocks for enhanced weathering (a CDR technique) was included in the top 20 (Table 2). This would cause a fundamental alteration of biogeochemical properties of the soil (pH, structure, etc.) with the potential to reduce soil biodiversity and disrupt the activity of the soil organisms that underpin overlying ecological communities (Jensen et al. 2003). An associated increase in the availability of nutrients could also feedback to alter the composition and productivity of plant communities (Dawson et al. 2012). The overall combined effects of changes to interdependent abiotic soil properties – such as temperature, physical structure and biogeochemistry – are difficult to predict (Davidson et al. 1998) and understanding of soil dynamics and biota, and their interactions with above-ground systems, requires more research (De Deyn & van der Putten 2005). Similar concerns were raised in relation to the application of biochar to soil as a means to increase carbon sequestration (another CDR technique), as the effects of this technique on soil biodiversity are poorly understood (Lehmann et al. 2011).

3.2. Priority areas for research

Five climate engineering techniques (Table 1) were found in existing assessments to have higher anticipated technical feasibility and efficacy than other techniques (e.g. The Royal Society 2009; Vaughan & Lenton 2011). Of the SRM techniques, stratospheric sulfate aerosols and enhanced marine cloud albedo are relatively well-studied through model simulations and inter-comparisons, and both anticipated to have high potential effectiveness in counteracting climate change (Kravitz, Caldeira et al. 2013). Of the CDR techniques, bioenergy with carbon capture and storage (BECCS) uses techniques that are already well developed (International Energy Agency 2011) and has good carbon sequestration potential (Caldeira et al. 2013). It is also included in mitigation scenarios in the recent IPCC Fifth Assessment report (van Vuuren et al. 2011; IPCC 2014). Ocean fertilization with iron is receiving ongoing commercial interest and field trials demonstrate that it is possible, even if its ability to absorb and store atmospheric carbon dioxide over the long-term appears to be low (Strong et al. 2009; Williamson et al. 2012). Direct air capture (DAC) was also found to be pertinent to consider as there is ongoing research and development of potential technology designs (e.g. Choi et al. 2011).

For each of these techniques, the index of priority was used to identify the highest priority environmental changes that they could cause if implemented. For each change, the expert group identified key knowledge gaps and research questions about its biodiversity and ecosystem effects, detailed in Table 3.

3.2.1. Reinforcing current research priorities

Many of the questions are relevant to existing research priorities in ecological science, but climate engineering presents an important and unique context for investigation. For example, ‘What are the rates of warming that species can tolerate by means of adaptation or migration ... ?’ (Table 3) is a key area of research in relation to climate change (e.g.

Table 3. Priority research questions relating to the highest priority environmental changes associated with each of the five shortlisted climate engineering techniques. The 'Index of priority' combines their importance score and scientific understanding score; environmental changes with high importance and low scientific understanding of the biodiversity and ecosystem consequences were considered priorities for research.

Technique	Prioritized environmental changes	Index of priority	Suggested priority research questions
1. Stratospheric sulfate aerosols	Termination effect: Rapid increase of global temperatures if solar radiation management fail or are terminated	89.9	1. What are the rates of warming that species can tolerate by means of adaptation or migration and which key species and ecosystem-level processes are most vulnerable to such rapid changes? 2. Does a rapid increase in temperature modify the effects of other important stressors, and what are the synergistic effects of these multiple stressors on biodiversity and ecosystems? 3. What consequences does an abrupt change from cooling to rapid warming have for evolutionary adaptation to warming? 1. What is the effect on primary productivity of the combined influence of increased CO ₂ concentrations and reduced temperatures for the dominant plant species in major terrestrial biomes and for oceanic phytoplankton? 2. How will enhanced CO ₂ concentrations and reduced global temperatures impact on ocean uptake of CO ₂ and acidification rates and what are the implications for calcifying organisms and their role in transferring particulate organic carbon to the deep ocean? 3. What are the indirect effects of high atmospheric CO ₂ levels and reduced temperature on biodiversity and ecosystem structure and function, including the effects on taxa other than primary producers and as a result of impacts cascading through food webs? 1. How will changes in precipitation affect aridification and regional distributions of species and communities, especially trophic levels other than primary producers, and what implications does this have for ecosystem processes they control? 2. What impacts do variations in precipitation regimes have on belowground processes, including water uptake and root structure, over the medium to long term? 3. In marine habitats, how might changes in freshwater inputs to the ocean affect the intensity and distribution of acidification in the marine surface layer and ocean interior, and how does this affect ocean biodiversity and ecosystem function in various regions?
2. Enhanced marine cloud albedo	Regionally-variable changes in precipitation due to altered atmospheric circulation. Increase in some areas, decrease in others.	75	<i>[Prioritized environmental changes for this technique are the same as for 1. Stratospheric sulfate aerosols – they are common to both]</i>
3. Ocean fertilization with iron	Increased primary productivity in high nutrient low chlorophyll regions of the ocean	66	1. What are the taxon-specific responses of phytoplankton to fertilization in terms of their growth and chemical composition (C, N, P, Si and Fe stoichiometry) under different states of nutrient (in)sufficiency, and how should these responses be included in models of community and ecosystem response? 2. What ecosystem effects might occur beyond the fertilization zone (e.g. through changes in downstream nutrient regimes, changes in flux to deeper ocean communities)?

(Continued)

Table 3. (Continued)

Technique	Prioritized environmental changes	Index of priority	Suggested priority research questions
4. Biofuels with carbon capture and storage (BECCS)	Increase in anoxic or hypoxic regions in mid and deep oceans due to increased respiration during decomposition of additional organic matter	55	3. How might higher trophic levels (including zooplankton, fish and mammals) respond to enhanced throughput of organic material, due to large-scale and long-term fertilization, and how might such effects influence areas beyond the fertilization zone? 1. What are the likely rates of biological degradation of the organic matter generated by iron fertilization in deep, cold ocean environments and would the character of the material (e.g. carbon:nitrogen ratio) make a difference to mineralization rates?
			2. What is the anticipated scale of the impact of substantially increased input of organic matter (and its subsequent decomposition) on mid-water oxygen levels; will existing oxygen minimum zones be expanded or new ones created?
			3. How might increased volumes of anoxic water directly or indirectly impact higher trophic levels, for example, fish and mammals (e.g. on geographical and depth ranges, migration routes, physiological processes, prey availability and foraging etc.)?
4. Biofuels with carbon capture and storage (BECCS)	Conversion of habitats to large-scale production of biofuel feedstocks	56	1. What strategies for feedstock production – in terms of location and size of production type of existing land-use or habitat replaced, and size and connectivity of remaining natural areas – could we use such that biodiversity and/or ecosystem service loss is minimized per unit energy produced for different biofuel types?
			2. Which management regimes used for planting, growing and harvesting each type of biofuel feedstock will have the smallest impact on biodiversity and ecosystem services?
			3. Which biofuel crops in which location will provide the most energy whilst having the least impact on biodiversity and ecosystem services per unit area, and how can we properly assess the trade-off between the value of biofuel production and the loss of biodiversity/ecosystem services?
5. Direct air capture (DAC)	Biodiversity and ecosystem impacts of species used in feedstocks (e.g. introduced fast-growing tree varieties, invasive species etc.)	52	1. Can structurally complex, multispecies biofuel plantations be established that have adequate biomass production for economic viability, whilst also providing habitat for native species and other non-biofuel ecosystem services?
			2. Is the long term net impact on biodiversity and ecosystem services less if a small area of highly productive, high water demanding, agrochemical dependent and potentially invasive biofuel crops is established, relative to the impact of developing a larger area for biofuels, which although less productive, are also less water-demanding, agrochemical dependent and less likely to become invasive?
			3. Which genetic and agronomic methods could be used to reduce the risk of invasiveness and the need for agrochemicals, whilst increasing productivity and water use efficiency of biofuel crops?
5. Direct air capture (DAC)	Construction of large air-capturing structures on open areas of land	33	1. Which locations could be most suitable for the placement of the DAC structures and what is the profile of the ecosystems and biodiversity that currently exist there? (i.e. are species rare/unique/endemic? How resilient are communities to disturbance?)

(Continued)

Table 3. (Continued)

Technique	Prioritized environmental changes	Index of priority	Suggested priority research questions
			<div>2. How large will the footprint of the DAC structures be and will they present an influential obstacle in the landscape, causing potential interference to species' feeding, nesting or migratory activity?</div> <div>3. To what degree will habitats be altered and disturbed by the construction and maintenance of direct air capture structures? (e.g. will land need to be cleared? Will permanent access routes be established and frequently used?)</div>
	Contamination of air 'downstream' of DAC if reactive chemicals used to capture CO ₂ evaporate	42	<div>1. Will the likely concentration of chemicals in air passing through the DAC structure represent a biologically-significant level to species in surrounding ecosystems?</div> <div>2. How far from direct air capture structures might species be impacted by air contamination effects?</div> <div>3. How will contamination impact species' fitness and the structure of communities in habitats where DAC structures are established?</div>

(Schloss et al. 2012; Quintero & Wiens 2013; Peck et al. 2014). It is also critical to consider within the context of climate engineering. Atmospheric and stratospheric SRM ('dimming') techniques will cause global-scale reduction in incoming radiation leading to stabilized or reduced rates of warming. With intensive implementation, abrupt termination of the techniques would be expected to cause a rapid rise in global mean temperatures – the 'termination effect' – unless additional actions had been used in the interim to reduce atmospheric CO₂ (Matthews & Caldeira 2007; Jones et al. 2013). Some of the ecological impacts of the termination effect can be anticipated from ongoing research into the effects of ongoing climate change which indicates that warming could alter species distributions, migration patterns, breeding etc. (Cotton 2003; Hurlbert 2012). However, the rate of temperature increase associated with the termination effect at intensive SRM implementation is likely to be much more rapid. Rates of change could exceed the ability of many species to adapt or migrate (Bellard et al. 2012; Cahill et al. 2013; Quintero & Wiens 2013) which could lead to local extinctions and substantial changes in community assemblages (Willis et al. 2010). Palaeoecological records suggest that global biodiversity showed resilience to similar rapid temperature changes during the last glacial-interglacial transition (Willis et al. 2010), but modern pressures including habitat fragmentation and degradation may now limit the capacity of species to track changes. Overall, there still remain large uncertainties about the exact nature of the ecological impacts of global temperature rises and scientific understanding of the biodiversity and ecosystem effects of the termination effect was judged by the group to be low (Table 3). The intensity of the effects could however be much less if a more moderate approach to SRM implementation was used. For example, if techniques were implemented at a scale to induce only a small degree of cooling (Kosugi 2013) or to curtail the rate of warming in parallel with emissions reduction efforts (MacMartin et al. 2014)

Similarly, several of the research questions identified in relation to BECCS (Table 3) are existing priority topics of research in relation to biofuels for energy (Gove et al. 2010; Fletcher et al. 2011; Wiens et al. 2011). Overall, the effects of biomass production were considered to be well understood compared to other environmental changes assessed (scores in Supporting Information S4). However, the significant scale of production required for BECCS as a climate engineering technique represents a significant additional demand for feedstocks, reinforcing the importance of research effort on the ecological effects of such production.

3.2.2. Novel research areas

Other environmental changes predicted to be caused by climate engineering create relatively novel conditions compared both to conditions observed in the past, and to projected trajectories of ongoing climate and environmental change. The ecological effects of these changes are relatively less well understood. For example, reduced incoming solar radiation caused by atmospheric and stratospheric SRM techniques will lead to reduced rates of global warming. However, in the absence of measures to address greenhouse gas emissions, atmospheric CO₂ levels would remain high. This high CO₂, low temperature climate differs from both current conditions and the high temperature, high CO₂ conditions projected under future emissions scenarios (Secretariat of the Convention on Biological Diversity 2012) and represents a relatively novel global climate compared to current, historical or paleo-historical conditions (Williams et al. 2007; Tilmes et al. 2013). Temperature and CO₂ control fundamental ecological processes and the relative influence of the two parameters is highly complex

(Long et al. 2004). Climate and vegetation models suggest that elevated CO₂ would be the dominant influence and could reduce water stress of plants leading to enhanced terrestrial primary productivity in almost all regions (Long et al. 2004; Wiens et al. 2011; Donohue et al. 2013), but there is a large degree of uncertainty in these projections (Jones et al. 2013; Kravitz, Caldeira et al. 2013). Individual species, functional groups and biomes will also vary in their response to temperature and CO₂ levels (Higgins & Scheiter 2012; De Frenne et al. 2013). The potential to predict these effects is currently limited by factors including the low-resolution representation of ecological interactions in integrated global scale models (Mustin et al. 2007; Ostle & Ward 2012). Scientific understanding of the effects was judged to be low (see Supporting Information S4).

Even when environmental changes have historical natural proxies, there often remain knowledge gaps about their biodiversity and ecosystem effects. For example, implications of increased primary productivity in high nutrient low chlorophyll ocean regions with iron fertilization can be anticipated to some extent from observations of natural fertilization from deep water upwelling (Blain et al. 2007) or deposition of air borne dust (Martinez-Garcia et al. 2014). However, the complexity of ocean systems and possible feedbacks mean that certainty about the ecological effects remains low, reflected in the expert group scientific understanding score (Table 3). Questions like 'What ecosystem effects might occur beyond the fertilization zone ... ?' would require dedicated investigation should this climate engineering technique be implemented.

The suggested research questions (Table 3) demonstrate critical knowledge gaps about ecological effects of climate engineering, which will need to be addressed if the techniques are pursued. Many relate to topics already recognized by the ecological research community as priority knowledge gaps, but in the climate engineering context, may require investigation over different scales, timeframes and locations. Others relate to novel conditions that could be created by climate engineering, which raise new questions about potential biodiversity and ecosystem impacts.

3.3. Concluding remarks

3.3.1. Inclusion of biodiversity and ecosystem effects in climate engineering research and decision-making

In the discussion about climate engineering to date, potential biodiversity and ecosystem impacts of the techniques have received little attention and there has been very limited work by the ecological research community on this topic. We believe it has thus far been challenging to identify discrete research questions due to the scale, number, range and complexity of potential biodiversity and ecosystem effects. In addition, there is perhaps reluctance to engage with climate engineering, given that it involves large-scale manipulation of the earth system and is viewed by some as a distraction from reducing greenhouse-gas emissions.

In an effort to encourage timely research into the biodiversity and ecosystem impacts of climate engineering, we have reviewed a comprehensive range of potential effects and made a critical first attempt to prioritize them based on assessment of the importance of their biodiversity and ecosystem effects and the degree of scientific understanding about them. In doing so, we have identified some key knowledge gaps and questions. Some of these fit within research priorities already identified by ecological science, but climate engineering presents a novel application and extension of the investigations and reinforces the need to

investigate these topics further. Others relate to conditions potentially created by climate engineering that differ from past conditions and from those projected under underlying climate and environmental change.

Discussions – and decisions – on the governance of climate engineering are already occurring, e.g. recent amendments to the London Protocol (International Maritime Organization 2013; Schafer et al. 2013). For sound policy decisions to be made, it is critical that they are based on good scientific understanding. We hope our identification of key knowledge gaps and suggested research questions will act as a platform for more detailed consideration of the ecological implications of climate engineering from now on, both from the ecological research community, and from those working on climate engineering and related policy.

3.3.2. Expert consultation and uncertainty

Expert elicitation can help enhance limited information available from scientific study (Martin et al. 2012). It is useful in the case of climate engineering as empirical studies of the techniques are logistically difficult or impossible to conduct at the scales necessary (Secretariat of the Convention on Biological Diversity 2012). Extrapolation from analogous natural processes (for example, global dimming caused by volcanic eruptions; Robock et al. 2013) and climate envelope modeling (Couce et al. 2013) can inform expectations of future scenarios to some extent (Robock et al. 2013), but are less effective when conditions will be novel relative to the past (Sutherland 2006).

The expert group used their collective knowledge to interpret available information to identify which biodiversity and ecosystem effects of climate engineering from a long and diverse list are important to investigate further. They acknowledged complexities of the potential ecological effects of climate engineering not previously acknowledged in the climate engineering literature. For example, the importance of distinguishing the effects of ocean fertilization with iron from those associated with nitrogen or phosphorus, and the need to particularly consider vulnerability of island biodiversity.

Inevitably, there are sources of uncertainty and variability inherent in expert consultation. Our outcomes may have been different with a different group of experts due to varying knowledge and opinion on the ecological impacts being discussed. Outcomes also depend very much on how the issues are framed, such as the context in which climate engineering is considered. For example, whilst it was specified that the working group should consider the effects against a background of a warming world with an acidifying ocean, it was left up to the individual to interpret whether that should be a 'business as usual' scenario or one with low, medium or high global mitigation effort. As noted in the introduction, we also did not consider the effects of the overall climate amelioration that would occur if climate engineering were effective, which would also have considerable biodiversity and ecosystem effects, including some likely benefits.

There are also many uncertainties related to climate engineering that make anticipating biodiversity and ecosystem effects challenging. Most technologies are in the early stages of design and it is difficult to predict how they might evolve. The location, timing and scale of any future deployment of such techniques are all theoretical (Keith 2000), making it difficult to identify the specific circumstances under which the environmental changes would occur (The Royal Society 2009; Russell et al. 2012). This significant topic of ongoing research should occur in parallel with attempts to project biodiversity and ecosystem effects of climate

engineering. Biodiversity experts and climate engineering impact modelers should collaborate in order to produce reasonable scenarios of deployment (Carey & Burgman 2008) (and see Cusack et al. 2014).

4. Conclusion

Any climate engineering technique designed to alter the global climate will have significant implications for biodiversity and ecosystems. This study makes a first attempt to identify effects related to currently-discussed techniques that are priorities for detailed investigation. The outcomes should be considered for what it is: an assessment by a group of experienced researchers based on currently available information. It is not an evaluation of the relative benefits or risks of climate engineering. It is a scoping of knowledge gaps and research priorities related to the biodiversity and ecosystem effects of implementing the techniques. The major themes identified show the types of ecological impacts that are particularly critical to consider, and highlight both important overlaps with existing research priorities and knowledge gaps that require new research focus. If interest in climate engineering continues, biodiversity and ecosystem consequences must be comprehensively considered so that unintended consequences are avoided and any potential co-benefits are realized. Further horizon scanning and expert consultation processes similar to those used here could be valuable in identifying emerging issues.

Authors and contributors

RS and SS conducted the initial literature review of climate engineering effects, with subsequent input from CGM, WB and PI. CGM and WJS designed the study process and delivered the workshop along with WB, PI and JJB. JJB contributed significantly to the literature review of the technical feasibility of climate engineering techniques. All other authors (except TA) completed the survey scoring task and attended the workshop. TA analyzed the output data. CGM wrote the first draft of the manuscript, and all authors contributed substantially to revisions. WJS, WB and PI in particular made significant contributions to the direction and content of the manuscript.

Acknowledgements

This work was funded by the Institute for Advanced Sustainability Studies, who we thank for their efficient organization and hospitality for the workshop. We also thank the various experts who completed the survey but could not attend the workshop: Tom Battin, Richard T. Conant, Jason Hall-Spencer, Sandra Lavorel and Klaus Lorenz. We are also grateful to the Cambridge Conservation Initiative (CCI) Shared Challenges Programme who funded the initial literature review, and to Rosamunde Almond (CCI) who was involved in the conception of the project. WJS is funded by Arcadia

Disclosure statement

No potential conflict of interest was reported by the authors.

References

- Anthony KRN, Kleypas JA, Gattuso, J-P. **2011**. Coral reefs modify their seawater carbon chemistry – implications for impacts of ocean acidification. *Glob Change Biol.* 17:3655–3666. doi:[10.1111/j.1365-2486.2011.02510.x](https://doi.org/10.1111/j.1365-2486.2011.02510.x).
- Bailey DM, Collins MA, Gordon JDM, Zuur AF, Priede IA. **2009**. Long-term changes in deep-water fish populations in the northeast Atlantic: a deeper reaching effect of fisheries? *Proc R Soc B.* 276:1965–1969.
- Bellard C, Bertelsmeier C, Leadley P, Thuiller W, Courchamp F. **2012**. Impacts of climate change on the future of biodiversity. *Ecol Lett.* 15:365–377.
- Biernacki P, Waldorf D. **1981**. Snowball Sampling: problems and techniques of chain referral sampling. *Socio Meth Res.* 10:141–153.
- Blain S, Queguiner B, Armand L, Belviso S, Bombled B, Bopp L, Bowie A, Brunet C, Brussaard C, Carlotti F, et al. **2007**. Effect of natural iron fertilization on carbon sequestration in the Southern Ocean. *Nature.* 446:1070–1074. Available from: http://www.nature.com/nature/journal/v446/n7139/supinfo/nature05700_S1.html
- Bopp L, Resplandy L, Orr JC, Doney SC, Dunne JP, Gehlen M, Halloran P, Heinze C, Ilyina T, Séférian R, et al. **2013**. Multiple stressors of ocean ecosystems in the 21st century: projections with CMIP5 models. *Biogeosciences.* 10:6225–6245. doi:[10.5194/bg-10-6225-2013](https://doi.org/10.5194/bg-10-6225-2013).
- Burrows MT, Schoeman DS, Buckley LB, Moore P, Poloczanska ES, Brander KL, Brown C, Bruno JF, Duarte CM, Halpern BSH, et al. **2011**. The pace of shifting climate in marine and terrestrial ecosystems. *Science.* 334:652–655.
- Cahill AE, Aiello-Lammens ME, Fisher-Reid MC, Hua X, Karanewsky CJ, Ryu HY, Sbeglia GC, Spagnolo F, Waldron JB, Warsi O, Wiens JJ. **2013**. How does climate change cause extinction? *Proc R Soc B.* 280:20121890. doi: <http://dx.doi.org/10.1098/rspb.2012.1890>.
- Caldeira K, Govindasamy B, Cao L. **2013**. The science of geoengineering. *Annu Rev Earth Planet Sci.* 41:231–256.
- Carey JM, Burgman MA. **2008**. Linguistic uncertainty in qualitative risk analysis and how to minimize it. *Ann NY Acad Sci.* 1128:13–17.
- Chen IC, Hill JK, Ohlemuller R, Roy DB, Thomas CD. **2011**. Rapid range shifts of species associated with high levels of climate warming. *Science.* 333:1024–1026.
- Choi S, Drese JH, Eisenberger PM, Jones CW. **2011**. Application of amine-tethered solid sorbents for direct CO₂ capture from the ambient air. *Environ Sci Tech.* 45:2420–2427.
- Cleland EE, Collins SL, Dickson TL, Farrer EC, Gross KL, Gherardi LA, Hallett LM, Hobbs RJ, Hsu JS, Turnbull L, Suding KN. **2013**. Sensitivity of grassland plant community composition to spatial vs. temporal variation in precipitation. *Ecology.* 94:1687–1696. doi:[10.1890/12-1006.1](https://doi.org/10.1890/12-1006.1).
- Costello MJ, Coll M, Danovaro R, Halpin P, Ojaveer H, Miloslavich P. **2010**. A census of marine biodiversity knowledge, resources, and future challenges. *PLoS ONE.* 5:e12110. doi:[10.1371/journal.pone.0012110](https://doi.org/10.1371/journal.pone.0012110).
- Cotton PA. **2003**. Avian migration phenology and global climate change. *Proc Nat Acad Sci.* 100:12219–12222.
- Couce E, Irvine PJ, Gregorie LJ, Ridgwell A, Hendy EJ. **2013**. Tropical coral reef habitat in a geoengineered, high-CO₂ world. *Geophys Res Lett.* 40:1799–1805. doi:[10.1002/grl.50340](https://doi.org/10.1002/grl.50340).
- Cusack DF, Aksen J, Shwom R, Hartzell-Nichols L, White S, Mackey KRM. **2014**. An interdisciplinary assessment of climate engineering strategies. *Front Ecol Environ.* 12:280–287. doi:[10.1890/130030](https://doi.org/10.1890/130030).
- Davidson EA, Belk E, Boone RD. **1998**. Soil water content and temperature as independent or confounded factors controlling soil respiration in a temperate mixed hardwood forest. *Glob Change Biol.* 4:217–227.
- Dawson W, Fischer M, van Kleunen M. **2012**. Common and rare plant species respond differently to fertilisation and competition, whether they are alien or native. *Ecol Lett.* 15:873–880.
- De Deyn GB, van der Putten W. **2005**. Linking aboveground and belowground diversity. *TREE.* 20:625–633.
- De Frenne P, Rodríguez-Sánchez F, Coomes DA, Baeten L, Verstraeten G, Vellend M, Bernhardt-Römermann M, Brown CD, Brunet J, Cornelis J, et al. **2013**. Microclimate moderates plant responses to macroclimate warming. *Proc Nat Acad Sci.* 110:18561–18565. doi:[10.1073/pnas.1311190110](https://doi.org/10.1073/pnas.1311190110).

- Devine JA, Baker KD, Haedrich RL. **2006**. Fisheries: deep-sea fishes qualify as endangered. *Nature*. 439:29.
- Donohue RJ, Roderick ML, McVicar, TR, Farquhar, GD. **2013**. Impact of CO₂ fertilization on maximum foliage cover across the globe's warm, dry environments. *Geophysical Research Letters*. 40:3031–3035.
- Drinkwater K, Hunt G, Lehodey P, Lluch-Cota S, Murphy EJ, Sakuri Y, Schwing F, Beaugrand G, Svein S. **2010**. Climate forcing on marine ecosystems; p. 11–36. In: Barange M, Field JG, Harris RP, Hofmann EE, Perry I, Werner F, editor. *Marine ecosystems and global change*: Oxford Scholarship Online. Oxford University Press: Oxford.
- Dutreuil S, Bopp L, Tagliabue A. **2009**. Impact of enhanced vertical mixing on marine biogeochemistry: lessons for geo-engineering and natural variability. *Biogeosciences*. 6:901–912. doi:10.5194/bg-6-901-2009.
- Edwards M, Richardson AJ. **2004**. Impact of climate change on marine pelagic phenology and trophic mismatch. *Nature*. 430:881–884. doi:10.1038/nature02808; Available from: http://www.nature.com/nature/journal/v430/n7002/supinfo/nature02808_S1.html
- Falkowski P, Barber RT, Smetacek V. **1998**. Biogeochemical controls and feedbacks on ocean primary production. *Science*. 281:200–206.
- Fletcher RJ, Robertson BA, Evans J, Doran PJ, Alavalapati JRR, Schemske DW. **2011**. Biodiversity conservation in the era of biofuels: risks and opportunities. *Front Ecol Environ*. 9:161–168.
- Foster J, Cooper G, Galbraith L. **2013**. Patent application: salt water spray systems for cloud brightening droplets and nano-particle generation. US patent application PCT/US2013/020589. Washington (DC): U.S. Patent and Trademark Office.
- Gove B, Flower KA, Bradbury RB. **2010**. A review of environmental consequences of biomass production for UK energy consumption. Sandy: RSPB.
- Hardman-Mountford NJ, Polimene L, Hirata T, Brewin RJ, Aiken J. **2013**. Impacts of light shading and nutrient enrichment geo-engineering approaches on the productivity of a stratified, oligotrophic ocean ecosystem. *J R Soc Interface*. 10:20130701.
- Hartmann J, West AJ, Renforth P, Köhler P, De La Rocha CL, Wolf-Gladrow DA, Dürr HH, Scheffran J. **2013**. Enhanced chemical weathering as a geoengineering strategy to reduce atmospheric carbon dioxide, supply nutrients, and mitigate ocean acidification. *Rev Geophys*. 51:113–149.
- Higgins SI, Scheiter S. **2012**. Atmospheric CO₂ forces abrupt vegetation shifts locally, but not globally. *Nature*. 488:209–212.
- Hoffmann AA, Sgro CM. **2011**. Climate change and evolutionary adaptation. *Nature*. 470:479–485.
- Holt J. **1996**. Balancing task allocation in teams. In: Robertson S, editor. *Contemporary ergonomics*. London: Taylor and Francis; p. 351.
- Hurlbert AHL. **2012**. Spatiotemporal variation in avian migration phenology: citizen science reveals effects of climate change. *PLoS ONE*. 7:e31662. doi: 10.1371/journal.pone.0031662.
- Hutchings A, Raine R. **2006**. A systematic review of factors affecting the judgments produced by formal consensus development methods in health care. *J Health Serv Res Policy*. 11:172–179.
- International Energy Agency. **2011**. Technology Roadmap: Biofuels for Transport. Paris: International Energy Agency.
- International Energy Agency. **2015**. Energy and Climate Change: World Energy Outlook Special Report. Paris: International Energy Agency.
- International Maritime Organization. **2013**. Marine geoengineering including ocean fertilization to be regulated under amendments to international treaty. [cited 2013 Nov 7] Available from: <http://www.imo.org/MediaCentre/PressBriefings/Pages/45-marine-geoengineering.aspx>
- IPCC. **2005**. IPCC special report on carbon dioxide capture and storage. In: Metz B, Davidson O, de Coninck H, Loos M, Meyer L, editor. Cambridge: Cambridge University Press.
- Metz B, Davidson, O, de Coninck H, Loo M, Meyer, L, editors. IPCC special report on carbon dioxide capture and storage. Cambridge: Cambridge University Press.
- IPCC. **2013**. Working Group I Contribution to the IPCC Fifth Assessment Report Climate Change 2013: The Physical Science Basis Summary for Policymakers. Cambridge: Cambridge University Press.
- IPCC. **2014**. Working Group III – Mitigation of Climate Change: Technical Summary. Cambridge: Cambridge University Press.

- Irvine PJ, Ridgwell A, Lunt DJ. **2010**. Assessing the regional disparities in geoengineering impacts. *Geophys Res Lett*. 37:L18702. doi:[10.1029/2010GL044447](https://doi.org/10.1029/2010GL044447)
- Irvine PJ, Ridgwell A, Lunt DJ. **2011**. Climatic impacts of surface albedo geoengineering. *J Geophys Res*. 116:D24112. doi:[10.1029/2011JD016281](https://doi.org/10.1029/2011JD016281).
- Jackson RB, Canadell JG, Le Quéré C, Andrew RM, Korsbakken JI, Peters GP, Nakicenovic N. **2016**. Reaching peak emissions. *Nat Clim Chang*. 6:7–9.
- Jensen KD, Beier C, Michelsen A, Emmett BA. **2003**. Effects of experimental drought on microbial processes in two temperate heathlands at contrasting water conditions. *Appl Soil Ecol*. 24:165–176. doi:[10.1016/S0929-1393\(03\)00091-X](https://doi.org/10.1016/S0929-1393(03)00091-X).
- Jones A, Haywood JM, Alterskjær K, Boucher O, Cole JNS, Curry CL, Irvine PJ, Ji D, Kravitz B, Egill Kristjánsson J, et al. **2013**. The impact of abrupt suspension of solar radiation management (termination effect) in experiment G2 of the Geoengineering Model Intercomparison Project (GeoMIP). *J Geophys Res: Atmospheres*. 118:9743–9752. doi:[10.1002/jgrd.50762](https://doi.org/10.1002/jgrd.50762).
- Keith DW. **2000**. Geoengineering the climate: history and prospect. *Ann Rev Energy Env*. 25:245–284.
- Keller DP, Feng EY, Oschlies A. **2014**. Potential climate engineering effectiveness and side effects during a high carbon dioxide-emission scenario. *Nat Commun*. 5: 3304. doi:[10.1038/ncomms4304](https://doi.org/10.1038/ncomms4304).
- Kirk JTO. **2011**. Light and photosynthesis in aquatic ecosystems. 3rd ed. Cambridge: Cambridge University Press.
- Kosugi T. **2013**. Fail-safe solar radiation management geoengineering. *Mitig Adapt Strat Glob Chang*. 18:1141–1166.
- Kravitz B, Caldeira K, Boucher O, Robock A, Rasch PJ, Alterskjær K, Karam DB, Cole JNS, Curry CL, Haywood JM, et al. **2013**. Climate model response from the Geoengineering Model Intercomparison Project (GeoMIP). *J Geophys Res: Atmos*. 118:8320–8332. doi:[10.1002/jgrd.50646](https://doi.org/10.1002/jgrd.50646).
- Kravitz B, Robock A, Irvine P. **2013**. Robust results from climate model simulations of geoengineering. *Eos, Trans Amer Geophys Union*. 94:292–292. doi:[10.1002/2013eo330005](https://doi.org/10.1002/2013eo330005).
- Kroeker KJ, Kordas RL, Crim R, Hendriks IE, Ramajo L, Singh GS, Duarte CM, Gattuso J-P. **2013**. Impacts of ocean acidification on marine organisms: quantifying sensitivities and interaction with warming. *Glob Change Biol*. 19:1884–1896. doi:[10.1111/gcb.12179](https://doi.org/10.1111/gcb.12179).
- Lampitt RS, Achterberg EP, Anderson TR, Hughes JA, Iglesias-Rodriguez MD, Kelly-Gerreyn BA, Lucas M, Popova EE, Sanders R, Shepherd JG, et al. **2008**. Ocean fertilization: a potential means of geoengineering? *Philos Trans R Soc A*. 366:3919–3945.
- Latham J, Bower K, Choularton T, Coe H, Connolly P, Cooper G, Craft T, Foster J, Gadian A, Galbraith L. **2012**. Marine cloud brightening. *Philos Trans R Soc A*. 370:4217–4262.
- Lehmann J, Rillig MC, Thies J, Masiello CA, Hockaday WC, Crowley D. **2011**. Biochar effects on soil biota – a review. *Soil Biol Biochem*. 43:1812–1836.
- Lenton TM, Vaughan NE. **2009**. The radiative forcing potential of different climate geoengineering options. *Atmos Chem Phys*. 9:5539–5561.
- Levin LA, Etter RJ, Rex MA, Gooday AJ, Smith CR, Pineda J, Stuart CT, Hessler RR, Pawson D. **2001**. Environmental influences on regional deep-sea species diversity. *Annu Rev Ecol Syst*. 32:51–93.
- Long SP, Ainsworth AE, Rogers A, Ort DR. **2004**. Rising atmospheric carbon dioxide: plants FACE the future. *Annu Rev Plant Biol*. 55:591–628.
- Loope L, Giambelluca T. **1998**. Vulnerability of island tropical montane cloud forests to climate change, with special reference to East Maui, Hawaii. *Clim Chang*. 39:503–517. doi:[10.1023/a:1005372118420](https://doi.org/10.1023/a:1005372118420).
- MacMartin DG, Caldeira K, Keith DW. **2014**. Solar engineering to limit the rate of temperature change. *Philos Trans R Soc A*. 372: 20140134. doi:[10.1098/rsta.2014.0134](https://doi.org/10.1098/rsta.2014.0134).
- MacMynowski DG, Keith DW, Caldeira K, Shin H-J. **2011**. Can we test geoengineering? *Energy Environ Sci*. 4:5044–5052.
- Martin TG, Burgman MA, Fidler F, Kuhnert PM, Low-Choy S, McBride M, Mengersen K. **2012**. Eliciting expert knowledge in conservation science. *Conserv Biol*. 26:29–38. doi:[10.1111/j.1523-1739.2011.01806.x](https://doi.org/10.1111/j.1523-1739.2011.01806.x).
- Martinez-Garcia A, Sigman DM, Ren H, Anderson RF, Straub M, Hodel DA, Jaccard SL, Eglinton TI, Haug GH. **2014**. Iron fertilization of the subantarctic ocean during the last ice age. *Science*. 343:1347–1350.
- Matthews HD, Caldeira K. **2007**. Transient climate-carbon simulations of planetary geoengineering. *Proc Nat Acad Sci*. 104:9949–9954.

- Matthews HD, Cao L, Caldeira K. 2009. Sensitivity of ocean acidification to geoengineered climate stabilisation. *Geophys Res Lett*. 36:L10809.
- Matthews S, O'Connor R, Plantinga AJ. 2002. Quantifying the impacts on biodiversity of policies for carbon sequestration in forests. *Ecol Econ*. 40:71–87. doi:[10.1016/S0921-8009\(01\)00269-5](https://doi.org/10.1016/S0921-8009(01)00269-5).
- McCusker K, Battisti DS, Bitz CM. 2012. The climate response to stratospheric sulfate injections and implications for addressing climate emergencies. *J Clim*. 25:3096–3116.
- Meehl GA. 1996. Vulnerability of freshwater resources to climate change in the tropical pacific region. *Water Air Soil Pollut*. 92:203–213. doi:[10.1007/bf00175566](https://doi.org/10.1007/bf00175566).
- Menzel A, Sparks TH, Estrella N, Koch E, Aasa A, Ahas R, Alm-Kübler K, Bissolli P, Braslavská OG, Briede A, et al. 2006. European phenological response to climate change matches the warming pattern. *Glob Change Biol*. 12:1969–1976. doi:[10.1111/j.1365-2486.2006.01193.x](https://doi.org/10.1111/j.1365-2486.2006.01193.x).
- Mineur F, Cook E, Minchin D, Bohn K, Macleod A, Maggs C. 2012. Changing coasts: marine aliens and artificial structures. *Oceanogr Mar Biol*. 50:189–234.
- Mustin K, Sutherland WJ, Gill JA. 2007. The complexity of predicting climate-induced ecological impacts. *Clim Res*. 35:165–175. doi:[10.3354/cr00723](https://doi.org/10.3354/cr00723).
- Ostle N, Ward S. 2012. Climate change and soil biotic carbon cycling soil ecology and ecosystem services. Oxford: Oxford University Press; p. 241–255.
- Peck LS, Morley SA, Richard JC. 2014. Acclimation and thermal tolerance in Antarctic marine ectotherms. *J Exp Biol*. 217:16–22.
- Peters GP, Marland G, Le Quéré C, Boden T, Canadell JG, Raupach MR. 2012. Rapid growth in CO₂ emissions after the 2008–2009 global financial crisis. *Nat Clim Chang*. 2:2–4. doi:[10.1038/nclimate1332](https://doi.org/10.1038/nclimate1332).
- Pörtner HO, Farrell AP. 2008. Physiol and climate change. *Science*. 322:690–692.
- Quintero I, Wiens JJ. 2013. Rates of projected climate change dramatically exceed past rates of climatic niche evolution among vertebrate species. *Ecol Lett*. 16:1095–1103.
- Ricke K, Morgan G, Allen M. 2010. Regional climate response to solar-radiation management. *Nat Geosci*. 3:537–541.
- Rickels W, Klepper G, Dovern J, Betz G, Brachatzek N, Cacean S, Gussow K, Heintzenberg J, Hiller S, Hoose C, et al. 2011. Large-scale intentional interventions into the climate system? Assessing the climate engineering debate scoping report conducted on behalf of the German Federal Ministry of Education and Research (BMBF). Kiel: Kiel Earth Institute.
- Robock A, MacMartin D, Duren R, Christensen M. 2013. Studying geoengineering with natural and anthropogenic analogs. *Clim Chang*. 121:445–458. doi:[10.1007/s10584-013-0777-5](https://doi.org/10.1007/s10584-013-0777-5).
- Russell LM, Rasch PJ, Mace GM, Jackson RB, Shepherd J, Liss P, Leinen M, Schimel D, Vaughan NE, Janetos AC. 2012. Ecosystem impacts of geoengineering: a review for developing a science plan. *Ambio*. 41:350–369. doi:[10.1007/s13280-13012-10258-13285](https://doi.org/10.1007/s13280-13012-10258-13285).
- Schafer S, Irvine PJ, Hubert A-M, Reichwein D, Low S, Stelzer H, Maas A, Lawrence MG. 2013. Field tests of solar climate engineering. *Nat Clim Chang*. 3:766–766. doi:[10.1038/nclimate1987](https://doi.org/10.1038/nclimate1987).
- Schloss CA, Nunez TA, Lawler JJ. 2012. Dispersal will limit ability of mammals to track climate change in the Western Hemisphere. *Proc Nat Acad Sci*. 109:8606–8611. doi:[10.1073/pnas1116791109](https://doi.org/10.1073/pnas1116791109).
- Secretariat of the Convention on Biological Diversity. 2012. Geoengineering in relation to the convention on biological diversity: technical and regulatory matters. CBD Technical Series No 66. Montreal; p. 152.
- Sims DW, Genner MJ, Southward AJ, Hawkins SJ. 2001. Timing of squid migration reflects North Atlantic climate variability. *Proc R Soc B*. 268:2067–2611.
- Singarayer JS, Ridgwell A, Irvine PJ. 2009. Assessing the benefits of crop albedo bio-geoengineering. *Environ Res Lett*. 4:045110. doi:[10.1088/1748-9326/4/4/045110](https://doi.org/10.1088/1748-9326/4/4/045110).
- Stahlschmidt ZR, DeNardo DF, Holland JN, Kotler BP, Kruse-Peebles M. 2011. Tolerance mechanisms in North American deserts: biological and societal approaches to climate change. *J Arid Environ*. 75:681–687.
- Steidl RJ, Litt AR, Matter WJ. 2013. Effects of plant invasions on wildlife in desert grasslands. *Wildlife Soc B*. 37:527–536. doi:[10.1002/wsb.308](https://doi.org/10.1002/wsb.308).
- Strong A, Chisholm S, Miller C, Cullen J. 2009. Ocean fertilization: time to move on. *Nature*. 461:347–348.
- Sutherland WJ. 2006. Predicting the ecological consequences of environmental change: a review of the methods. *J Appl Ecol*. 43:599–616. doi:[10.1111/j.1365-2664.2006.01182.x](https://doi.org/10.1111/j.1365-2664.2006.01182.x).

- Sutherland WJ, Bardsley S, Clout M, Depledge MH, Dicks LV, Fellman L, Fleishman E, Gibbons DW, Keim B, Lickorish F, et al. **2012**. A horizon scan of global conservation issues for 2013. *TREE*. 28:16–22.
- Sutherland WJ, Woodroof HJ. **2009**. The need for environmental horizon scanning. *TREE*. 24:523–527. doi:[10.1016/j.tree.2009.04.008](https://doi.org/10.1016/j.tree.2009.04.008).
- The Royal Society. **2009**. Geoengineering the climate – science, governance and uncertainty. RS Policy document 10/09. London: The Royal Society.
- Tilman D, Socolow R, Foley JA, Hill J, Larson E, Lynd L, Pacala S, Reilly J, Searchinger T, Somerville C, Williams R. **2009**. Beneficial biofuels – the food, energy, and environment trilemma. *Science*. 325:270–271. doi:[10.1126/science.1177970](https://doi.org/10.1126/science.1177970).
- Tilmes S, Fasullo J, Lamarque J-F, Marsh DR, Mills M, Alterskjær K, Muri H, Kristjánsson JE, Boucher O, Schulz M, et al. **2013**. The hydrological impact of geoengineering in the Geoengineering Model Intercomparison Project (GeoMIP). *J Geophys Res: Atmos*. 118:11036–11058. doi:[10.1002/jgrd.50868](https://doi.org/10.1002/jgrd.50868).
- van Vuuren DP, Edmonds J, Kainuma M, Riahi K, Thomson A, Hibbard K, Hurtt GC, Kram T, Krey V, Lamarque J-F, et al. **2011**. The representative concentration pathways: an overview. *Clim Chang*. doi:[10.1007/s10584-011-0148-z](https://doi.org/10.1007/s10584-011-0148-z).
- Vaughan NE, Lenton TM. **2011**. A review of climate geoengineering proposals. *Clim Chang*. 109:745–790.
- Walther G-R, Post E, Convey P, Menzel A, Parmesan C, Beebee TJC, Fromentin J-M, Hoegh-Guldberg O, Bairlein F. **2002**. Ecological responses to recent climate change. *Nature*. 416:389–395.
- Warren R, VanDerWal J, Price J, Walbergen JA, Atkinson I, Ramirez-Villegas J, Osborn TJ, Jarvis A, Shoo LP, Williams SE, Lowe J. **2013**. Quantifying the benefit of early climate change mitigation in avoiding biodiversity loss. *Nat Clim Chang*. 3:678–682.
- Wiens JJ, Fargione J, Hill J. **2011**. Biofuels and biodiversity. *Ecol Appl*. 21:1085–1095.
- Williams JW, Jackson ST, Kutzbach JE. **2007**. Projected distributions of novel and disappearing climates by 2100 AD. *Proc Nat Acad Sci*. 104:5738–5742. doi:[10.1073/pnas.0606292104](https://doi.org/10.1073/pnas.0606292104).
- Williamson P, Turley C. **2012**. Ocean acidification in a geoengineering context. *Philos Trans R Soc A*. 370:4317–4342. doi:[10.1098/rsta.2012.0167](https://doi.org/10.1098/rsta.2012.0167).
- Williamson P, Wallace DWR, Law CS, Boyd PW, Collos Y, Croot P, Denman K, Riebesell U, Takeda S, Vivian C. **2012**. Ocean fertilisation for geoengineering: a review of effectiveness, environmental impacts and emerging governance. *Process Saf Environ*. 90:475–488.
- Willis KJ, Bailey RM, Bhagwat SA, Birks HJB. **2010**. Biodiversity baselines, thresholds and resilience: testing predictions and assumptions using palaeological data. *TREE*. 25:583–591.
- Willis KJ, Bennett KD, Burrough SL, Macias-Fauria M, Tovar C. **2013**. Determining the response of African biota to climate change: using the past to model the future. *Philos Trans R Soc B: Biol Sci*. 368:20120491. doi: [10.1098/rstb.2012.0491](https://doi.org/10.1098/rstb.2012.0491).
- Winder MS. **2004**. Climate change uncouples trophic interactions in an aquatic ecosystem. *Ecology*. 85:2100–2106.
- Zhou S, Flynn PC. **2005**. Geoengineering downwelling ocean currents: a cost assessment. *Clim Chang*. 71:203–220.